

# MAT 425: Homework 11 (04/20)

**Ex 1.** Let  $\ell = 10$  and  $\alpha = .5$  and consider the parabolic PDE with **periodic boundary conditions**:

$$\begin{cases} \partial_t u = \alpha^2 \partial_x^2 u & , 0 < x < \ell, t > 0 \\ u(x, 0) = \sin \frac{\pi}{\ell} x & , 0 < x < \ell \\ \mathbf{u}(\mathbf{0}, \mathbf{t}) = \mathbf{u}(\ell, \mathbf{t}) & , t \geq 0. \end{cases}$$

- a) Implement a finite-difference method to solve this PDE.
- b) Consider the total mass of the solution:

$$m(t) = \int_0^\ell u(x, t) dx.$$

- Show (analytically) that the mass  $m(t)$  is conserved.  
*Hint: by periodicity  $\partial_x u(0, t) = \partial_x u(\ell, t)$  for all  $t \geq 0$ .*
  - Plot the evolution of the mass  $m(t)$  in time for the numerical solution.
- c) Suppose we use as initial condition:  $u_0(x) = 1 + \sin \frac{2\pi x}{\ell}$ .  
 Can you guess what will be the stationary state? (i.e. find  $u_*(x)$  such that  $u(x, t) \xrightarrow{t \rightarrow \infty} u_*(x)$ ).  
 Confirm numerically your guess.

**Ex 2.** Let  $\ell = 1$  and  $c = 1$  and consider the wave equation:

$$\begin{cases} \partial_t^2 u = c^2 \partial_x^2 u & , 0 < x < \ell, t > 0 \\ u(x, 0) = \sin 2\pi x & , 0 < x < \ell \\ \partial_t u(x, 0) = 2\pi \sin 2\pi x & , 0 < x < \ell \\ \mathbf{u}(\mathbf{0}, \mathbf{t}) = \mathbf{u}(\ell, \mathbf{t}) & , t \geq 0. \end{cases}$$

- a) Implement a finite-difference method to solve this PDE.
- b) Test the scheme with:

$$i) \Delta x = 2 \cdot 10^{-1}, \Delta t = 10^{-1} \quad , \quad ii) \Delta x = 10^{-1}, \Delta t = 2 \cdot 10^{-1}.$$

What do you observe?

- c) Find the accuracy of the scheme using  $\Delta t = .9 \cdot \Delta x$ .  
*Hint: take  $\Delta x = \frac{1}{10}, \frac{1}{30}, \frac{1}{70}, \frac{1}{200}, \frac{1}{550}, \frac{1}{1500}$  and compute the error with the exact solution  $u(x, t) = \sin 2\pi x \cdot (\cos 2\pi t + \sin 2\pi t)$ .*